ELSEVIER

Neuroscience Letters 351 (2003) 210–213

Neuroscience Letters

www.elsevier.com/locate/neulet

Author Index to Volume 351

Akema, T., see Maruki, K., 95
Albo, F., see Pieri, M., 153
Alisky, J.M., Hughes, S.M., Davidson, B.L., Transduction of neurons lining the cerebral external capsules in mice with feline immunodeficiency virus based vectors, 120
Almaguer-Melian, W., see Bergado, J.A., 56
Annesi, F., see Spadafora, P., 75
Annesi, G., see Spadafora, P., 75
Anschel, D.J., Pascual-Leone, A., Holmes, G.L., Anti-kindling effect of slow repetitive transcranial magnetic stimulation in rats, 9
Averill, S., see Delcroix, J.-D., 181

Barry, P.H., see Lee, D.J.-S., 196
Bengtson, C.P., see Pieri, M., 153
Berg, D., see Krüger, R., 125
Bergado, J.A., Almaguer-Melian, W., Kostenko, S., Frey, S., Frey, J.U., Behavioral reinforcement of long-term potentiation in rat dentate gyrus in vivo is protein synthesis-dependent, 56
Berger, K., see Krüger, R., 125
Bickford, M.E., see Carden, W.B., 87
Bohotin, C., Scholsem, M., Bohotin, V., Franzen, R., Schoenen, J., Vagus nerve stimulation attenuates heat- and formalin-induced pain in rats, 79
Bohotin, V., see Bohotin, C., 79
Borsdal, J., see Torup, L., 173
Bromberg, Y., see Sperling, O., 137

Carden, W.B., Guido, W., Godwin, D.W., Bickford, M.E., Thalamocortical cells in the cat pulvinar nucleus transiently express nitric oxide synthase during development, 87
Carrideo, S., see Spadafora, P., 75
Carter, S.J., Richardson, C.M., Wakerley, J.B., Excitatory effects of oxytocin and cholecystokinin on oxytocin neurones: differences between virgin, pregnant and lactating rats, 13
Cavalcanti, S., see Pieri, M., 153
Chen, K., Zhu, J.-N., Li, H.-Z., Wang, J.-J., Histamine elicits neuronal excitatory response of red nucleus in the rat via H_2 receptors in vitro, 25
Chen, R., Osterhaus, G., McKerchar, T., Fowler, S.C., The role of exogenous testosterone in cocaine-induced behavioral sensitization and plasmalemmal or vesicular dopamine uptake in castrated rats, 161
Chen, X.-Y., see Wang, Q., 83
Choi, S.-S., see Lee, H.-K., 209
Cirò Candiano, I.C., see Spadafora, P., 75
Civitelli, D., see Spadafora, P., 75
Cooper, J.M., see Orth, M., 29
Czepiel, K., see Przegaliński, E., 169

Davidson, B.L., see Alisky, J.M., 120
De Marco, E.V., see Spadafora, P., 75
Delcroix, J.-D., Patel, J., Averill, S., Tomlinson, D.R., Priestley, J.V., Fernyhough, P., Peripheral axon crush elevates transport of $p75^{NTR}$ in the central projection of sensory neurones of rats, 181
Dheen, S.T., see Lian, Q., 51
Djavadian, R.L., Wielkopolska, E., Turlejski, K., Neonatal depletion of serotonin increases the numbers of callosally projecting neurons in cat visual areas 17 and 18, 91
Dlaboga, D., see Przegaliński, E., 169
Dravid, S.M., Murray, T.F., Fluorescent detection of Ca^{2+}-permeable AMPA/kainate receptor activation in murine neocortical neurons, 145

Eguchi, Y., Ogiue-Ikeda, M., Ueno, S., Control of orientation of rat Schwann cells using an 8-T static magnetic field, 130
Epplen, J.T., see Krüger, R., 125

Fernyhough, P., see Delcroix, J.-D., 181
Filip, M.s., see Przegaliński, E., 169
Fischer, C., see Krüger, R., 125
Fowler, S.C., see Chen, R., 161
Franzen, R., see Bohotin, C., 79
Frey, J.U., see Bergado, J.A., 56
Frey, S., see Bergado, J.A., 56

Gaetti, C., see Pieri, M., 153
Gaudreau, G.-A., Plourde, V., Role of tachykinin NK_1, NK_2 and NK_3 receptors in the modulation of visceral hypersensitivity in the rat, 59
Gaus, S., see Suyama, H., 133
Gear, R.W., Gordon, N.C., Miaskowski, C., Paul, S.M., Heller, P.H., Levine, J.D., Dose ratio is important in maximizing naloxone enhancement of nalbuphine analgesia in humans, 5
Gerner, H.J., see Pfurtscheller, G., 33
Gobbele, R., see Krüger, R., 125
Godwin, D.W., see Carden, W.B., 87
Gong, Z.-H., see Yang, H.-J., 206
Gordon, N.C., see Gear, R.W., 5
Goshima, F., see Mori, I., 201
Gu, B., Zhang, Y.-D., Hu, G., A unilateral 6-hydroxydopamine lesion decreases the expression of metabotropic glutamate receptors in rat substantia nigra, 186
Guido, W., see Carden, W.B., 87

Höpfner, M., see Nguyen, D., 71
Hagan, M., Wennersten, A., Meijer, X., Holmin, S., Wahlberg, L., Mathiesen, T., Neuroprotection by human neural progenitor cells after experimental contusion in rats, 149
Haji, A., Yamazaki, H., Ohi, Y., Takeda, R., Distribution of μ receptors in the ventral respiratory group neurons; immunohistochemical and pharmacological studies in decerebrate cats, 37

doi:10.1016/S0304-3940(03)01202-3

Han, E.-J., see Lee, H.-K., 209
Han, H.C., see Zhang, G.H., 177
Han, K.-J., see Lee, H.-K., 209
Hasegawa, Y., see Saito, D., 107
Hashimoto, K., see Shimizu, E., 111
Hayes, E.A., Tiippana, K., Nicol, T.G., Sams, M., Kraus, N., Integration of heard and seen speech: a factor in learning disabilities in children, 46
Heller, P.H., see Gear, R.W., 5
Henke, U., see Nguyen, D., 71
Herness, S., see Lu, S.-g., 157
Holmes, G.L., see Anschel, D.J., 9
Holmin, S., see Hagan, M., 149
Honda, M., see Yagi, R., 191
Hong, S.K., see Zhang, G.H., 177
Howe, C.L., Depolarization of PC12 cells induces neurite outgrowth and enhances nerve growth factor-induced neurite outgrowth in rats, 41
Hu, G., see Gu, B., 186
Hughes, S.M., see Alisky, J.M., 120
Hungs, M., see Krüger, R., 125

Ishida, N., see Oishi, K., 21
Isomura, Y., see Tsujimoto, T., 103
Iyo, M., see Shimizu, E., 111
Izaki, Y., see Maruki, K., 95

Jones, T.A., see Sorensen, S.A., 17

Kawamoto, M., see Suyama, H., 133
Keramidas, A., see Lee, D.J.-S., 196
Kimura, Y., see Mori, I., 201
Koide, N., see Mori, I., 201
Koike, K., see Shimizu, E., 111
Komatsu, N., see Shimizu, E., 111
Koshizuka, T., see Mori, I., 201
Kostenko, S., see Bergado, J.A., 56
Krüger, R., Fischer, C., Schulte, T., Strauss, K.M., Müller, T., Woitalla, D., Berg, D., Hungs, M., Gobbele, R., Berger, K., Epplen, J.T., Riess, O., Schöls, L., Mutation analysis of the neurofilament M gene in Parkinson's disease, 125
Kraus, N., see Hayes, E.A., 46
Kreuder, F., see Westerhausen, R., 99
Kumakiri, C., see Shimizu, E., 111

Lee, D.J.-S., Keramidas, A., Moorhouse, A.J., Schofield, P.R., Barry, P.H., The contribution of proline 250 (P-2′) to pore diameter and ion selectivity in the human glycine receptor channel, 196
Lee, H.-K., Choi, S.-S., Han, E.-J., Han, K.-J., Suh, H.-W., Erratum to "Role of glutamate receptors and an on-going protein synthesis in the regulation of phosphorylation of Ca^{2+}/calmodulin-dependent protein kinase II in the CA3 hippocampal region in mice administered with kainic acid intracerebroventricularly" [Neuroscience Letters 348 (2003) 93–96], 209
Lee, K.S., see Zhang, G.H., 177
Levine, J.D., see Gear, R.W., 5
Li, H., see Wang, Q., 83
Li, H.-Z., see Chen, K., 25
Li, Y.-F., see Yang, H.-J., 206
Lian, Q., Samuel, T.S.W., Dheen, S.T., Enhanced expression of transforming growth factor-beta isoforms in the neural tube of embryos derived from diabetic mice exposed to cyclophosphamide, 51
Liu, J., see Wang, Q., 83
Llewellyn-Smith, I.J., Martin, C.L., Marcus, J.N., Yanagisawa, M., Minson, J.B., Scammell, T.E., Orexin-immunoreactive inputs to rat sympathetic preganglionic neurons, 115
Longone, P., see Pieri, M., 153
Lu, S.-g., Zhao, F.-l., Herness, S., Physiological phenotyping of cholecystokinin-responsive rat taste receptor cells, 157
Luo, Z.-P., see Yang, H.-J., 206

Müller, G.R., see Pfurtscheller, G., 33
Müller, T., see Krüger, R., 125
Marcus, J.N., see Llewellyn-Smith, I.J., 115
Martin, C.L., see Llewellyn-Smith, I.J., 115
Maruki, K., Izaki, Y., Akema, T., Nomura, M., Effects of acetylcholine antagonist injection into the prefrontal cortex on the progress of lever-press extinction in rats, 95
Mathiesen, T., see Hagan, M., 149
Maton, B., see Wicart, P., 67
McKerchar, T., see Chen, R., 161
Meijer, X., see Hagan, M., 149
Miaskowski, C., see Gear, R.W., 5
Min, S.S., see Zhang, G.H., 177
Minson, J.B., see Llewellyn-Smith, I.J., 115
Moorhouse, A.J., see Lee, D.J.-S., 196
Mori, I., Goshima, F., Koshizuka, T., Koide, N., Sugiyama, T., Yoshida, T., Yokochi, T., Kimura, Y., Nishiyama, Y., The US3 protein kinase of herpes simplex virus attenuates the activation of the c-Jun N-terminal protein kinase signal transduction pathway in infected piriform cortex neurons of C57BL/6 mice, 201
Murray, T.F., see Dravid, S.M., 145
Nakazato, M., see Shimizu, E., 111

Newman, W.H., see Wang, Z., 1
Nguyen, D., Höpfner, M., Zobel, F., Henke, U., Scherübl, H., Stangel, M., Rat oligodendroglial cell lines express a functional receptor for the chemokine CCL3 (macrophage inflammatory protein-1alpha), 71
Nicol, T.G., see Hayes, E.A., 46
Nicoletti, G., see Spadafora, P., 75
Nishina, E., see Yagi, R., 191
Nishiyama, Y., see Mori, I., 201
Nomura, M., see Maruki, K., 95
Nowak, E., see Przegaliński, E., 169

Oelsner, H., see Sperling, O., 137
Ogiue-Ikeda, M., see Eguchi, Y., 130
Ohi, Y., see Haji, A., 37
Oishi, K., Sakamoto, K., Ishida, N., Bimodal circadian expression of serotonin *N*-acetyltransferase mRNA in the retina of rats under restricted feeding, 21
Okada, S.-i., see Shimizu, E., 111
Okamura, N., see Shimizu, E., 111
Olavarria, J.F., see Sorensen, S.A., 17
Oohashi, T., see Yagi, R., 191
Orth, M., Tabrizi, S.J., Schapira, A.H.V., Cooper, J.M., α-Synuclein expression in HEK293 cells enhances the mitochondrial sensitivity to rotenone, 29
Osterhaus, G., see Chen, R., 161

Park, J.Y., see Zhang, G.H., 177
Pascual-Leone, A., see Anschel, D.J., 9

Pasqua, A.A., see Spadafora, P., 75
Patel, J., see Delcroix, J.-D., 181
Paul, S.M., see Gear, R.W., 5
Pfurtscheller, G., Müller, G.R., Pfurtscheller, J., Gerner, H.J., Rupp, R., 'Thought' – control of functional electrical stimulation to restore hand grasp in a patient with tetraplegia, 33
Pfurtscheller, J., see Pfurtscheller, G., 33
Pieri, M., Albo, F., Gaetti, C., Spalloni, A., Bengtson, C.P., Longone, P., Cavalcanti, S., Zona, C., Altered excitability of motor neurons in a transgenic mouse model of familial amyotrophic lateral sclerosis, 153
Plourde, V., see Gaudreau, G.-A., 59
Priestley, J.V., see Delcroix, J.-D., 181
Przegaliński, E., Czepiel, K., Nowak, E., Dlaboga, D., Filip, M.s., Withdrawal from chronic cocaine up-regulates 5-HT_{1B} receptors in the rat brain, 169

Quattrone, A., see Spadafora, P., 75

Rao, P.J., see Wang, Z., 1
Richardson, C.M., see Carter, S.J., 13
Riess, O., see Krüger, R., 125
Ruan, D.-Y., see Zhang, M.-M., 165
Rupp, R., see Pfurtscheller, G., 33

Sager, T., see Torup, L., 173
Saito, D., Hasegawa, Y., Urano, A., Gonadotropin-releasing hormones modulate electrical activity of vasotocin and isotocin neurons in the brain of rainbow trout, 107
Sakamoto, K., see Oishi, K., 21
Sams, M., see Hayes, E.A., 46
Samuel, T.S.W., see Lian, Q., 51
Sasaki, K., see Tsujimoto, T., 103
Scammell, T.E., see Llewellyn-Smith, I.J., 115
Schöls, L., see Krüger, R., 125
Schapira, A.H.V., see Orth, M., 29
Scherübl, H., see Nguyen, D., 71
Schoenen, J., see Bohotin, C., 79
Schofield, P.R., see Lee, D.J.-S., 196
Scholsem, M., see Bohotin, C., 79
Schulte, T., see Krüger, R., 125
Schweiger, E., see Westerhausen, R., 99
Segade, L.A.G., Trigeminal primary afferent projections to the peribrachial area in the guinea pig, 63
Serra, P., see Spadafora, P., 75
Shillcutt, S.D., see Wang, Z., 1
Shimazu, H., see Tsujimoto, T., 103
Shimizu, E., Hashimoto, K., Watanabe, H., Komatsu, N., Okamura, N., Koike, K., Shinoda, N., Nakazato, M., Kumakiri, C., Okada, S.-i., Iyo, M., Serum brain-derived neurotrophic factor (BDNF) levels in schizophrenia are indistinguishable from controls, 111
Shinoda, N., see Shimizu, E., 111
Sorensen, S.A., Jones, T.A., Olavarria, J.F., Neonatal enucleation reduces the proportion of callosal boutons forming multiple synaptic contacts in rat striate cortex, 17
Spadafora, P., Annesi, G., Pasqua, A.A., Serra, P., Cirò Candiano, I.C., Carrideo, S., Tarantino, P., Civitelli, D., De Marco, E.V., Nicoletti, G., Annesi, F., Quattrone, A., NACP-REP1 polymorphism is not involved in Parkinson's disease: a case-control study in a population sample from southern Italy, 75
Spalloni, A., see Pieri, M., 153
Sperling, O., Bromberg, Y., Oelsner, H., Zoref-Shani, E., Reactive oxygen species play an important role in iodoacetate-induced neurotoxicity in primary rat neuronal cultures and in differentiated PC12 cells, 137
Sreng, L., Sensory asymmetries in the olfactory system underlie sexual pheromone communication in the cockroach *Nauphoeta cinerea*, 141
Stangel, M., see Nguyen, D., 71
Strauss, K.M., see Krüger, R., 125
Sugiyama, T., see Mori, I., 201
Suh, H.-W., see Lee, H.-K., 209
Suyama, H., Kawamoto, M., Gaus, S., Yuge, O., Effect of JTC-801 (nociceptin antagonist) on neuropathic pain in a rat model, 133

Tabrizi, S.J., see Orth, M., 29
Takeda, R., see Haji, A., 37
Tarantino, P., see Spadafora, P., 75
Tiippana, K., see Hayes, E.A., 46
Tomlinson, D.R., see Delcroix, J.-D., 181
Torup, L., Borsdal, J., Sager, T., Neuroprotective effect of the neurotensin analogue JMV-449 in a mouse model of permanent middle cerebral ischaemia, 173
Tsujimoto, T., Shimazu, H., Isomura, Y., Sasaki, K., Prefrontal theta oscillations associated with hand movements triggered by warning and imperative stimuli in the monkey, 103
Turlejski, K., see Djavadian, R.L., 91

Ueno, S., see Eguchi, Y., 130
Urano, A., see Saito, D., 107

Wahlberg, L., see Hagan, M., 149
Wakerley, J.B., see Carter, S.J., 13
Walter, C., see Westerhausen, R., 99
Wang, J.-J., see Chen, K., 25
Wang, Q., Li, H., Wang, X.-W., Wu, D.-c., Chen, X.-Y., Liu, J., Resveratrol promotes differentiation and induces Fas-independent apoptosis of human medulloblastoma cells, 83
Wang, X.-W., see Wang, Q., 83
Wang, Z., Rao, P.J., Shillcutt, S.D., Newman, W.H., Phenotypic diversity of smooth muscle cells isolated from human intracranial basilar artery, 1
Watanabe, H., see Shimizu, E., 111
Wennersten, A., see Hagan, M., 149
Westerhausen, R., Walter, C., Kreuder, F., Wittling, R.A., Schweiger, E., Wittling, W., The influence of handedness and gender on the microstructure of the human corpus callosum: a diffusion-tensor magnetic resonance imaging study, 99
Wicart, P., Maton, B., Body equilibrium at the end of gait initiation: importance of ankle muscular force as evidenced in clubfoot children, 67
Wielkopolska, E., see Djavadian, R.L., 91
Wittling, R.A., see Westerhausen, R., 99
Wittling, W., see Westerhausen, R., 99
Woitalla, D., see Krüger, R., 125
Wu, D.-c., see Wang, Q., 83

Xiao, C., see Zhang, M.-M., 165

Yagi, R., Nishina, E., Honda, M., Oohashi, T., Modulatory effect of inaudible high-frequency sounds on human acoustic perception, 191
Yamazaki, H., see Haji, A., 37
Yanagisawa, M., see Llewellyn-Smith, I.J., 115
Yang, H.-J., Li, Y.-F., Zhang, H.-T., Zhang, F.-Q., Zhao, N., Gong, Z.-H., Luo, Z.-P., Up-regulation of microtubule-associated protein 4 and dreb-

rin A mRNA expression by antidepressants in rat hippocampus following chronic stress, 206
Yokochi, T., see Mori, I., 201
Yoon, Y.W., see Zhang, G.H., 177
Yoshida, T., see Mori, I., 201
Yu, K., see Zhang, M.-M., 165
Yuge, O., see Suyama, H., 133

Zhang, F.-Q., see Yang, H.-J., 206
Zhang, G.H., Yoon, Y.W., Lee, K.S., Min, S.S., Hong, S.K., Park, J.Y., Han, H.C., The glutamatergic *N*-methyl-D-aspartate and non-*N*-methyl-D-aspartate receptors in the joint contribute to the induction, but not maintenance, of arthritic pain in rats, 177
Zhang, H.-T., see Yang, H.-J., 206
Zhang, M.-M., Yu, K., Xiao, C., Ruan, D.-Y., The influence of developmental periods of sodium valproate exposure on synaptic plasticity in the CA1 region of rat hippocampus, 165
Zhang, Y.-D., see Gu, B., 186
Zhao, F.-l., see Lu, S.-g., 157
Zhao, N., see Yang, H.-J., 206
Zhu, J.-N., see Chen, K., 25
Zobel, F., see Nguyen, D., 71
Zona, C., see Pieri, M., 153
Zoref-Shani, E., see Sperling, O., 137